

The two INTEGRAL X-ray transients IGR J17091–3624 and IGR J17098–3628: a multi-wavelength long term campaign

F. Capitanio¹, M. Giroletti², M. Molina³, A. Bazzano¹, A. Tarana¹, J. Kennea⁴, A. J. Dean³, A. B. Hill³, M. Tavani¹, P. Ubertini¹

ABSTRACT

IGR J17091–3624 and IGR J17098–3628 are two X-ray transients discovered by *INTEGRAL* and classified as possible black hole candidates (BHCs). We present here the results obtained from the analysis of multi-wavelength data sets collected by different instruments from 2005 until the end of 2007 on both sources. IGR J17098–3628 has been regularly detected by *INTEGRAL* and *RXTE* over the entire period of the observational campaign; it was also observed with pointed observations by *XMM* and *Swift*/XRT in 2005 and 2006 and exhibited flux variations not linked with the change of any particular spectral features. IGR J17091–3624 was initially in quiescence (after a period of activity between 2003 April and 2004 April) and it was then detected again in outburst in the XRT field of view during a *Swift* observation of IGR J17098–3628 on 2007 July 9. The observations during quiescence provide an upper limit to the 0.2–10 keV luminosity, while the observations in outburst cover the transition from the hard to the soft state. Moreover, we obtain a refined X-ray position for IGR J17091–3624 from the *Swift*/XRT observations during the outburst in 2007. The new position is inconsistent with the previously proposed radio counterpart. We identify in VLA archive data a compact radio source consistent with the new X-ray position and propose it as the radio counterpart of the X-ray transient.

Subject headings: X-rays: binaries – X-rays: stars – X-rays: individual: IGR J17098–3628 – IGR J17091–3624 – black hole candidate

¹INAF IASF-Roma, Via Fosso del Cavaliere 100, 00033 Rome, Italy

²INAF Istituto di Radioastronomia, via Gobetti 101, 40129 Bologna, Italy

³School of Physics and Astronomy, University of Southampton, Highfield Southampton, SO17 1BJ, UK

⁴Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, Pennsylvania, USA

1. Introduction

Among the sources included in the *INTEGRAL* IBIS/ISGRI survey catalog (Bird et al. 2007), the two X-ray transients IGR J17091–3624 and IGR J17098–3628 are remarkable because of their proximity, being only 9'.6 away from each other. The two sources have been detected in different periods of time from April 2003 until the beginning of 2008. In particular, IGR J17091–3624 was detected by IBIS in 2003 and remained detectable for one year and, after a period of quiescence, it was again in outburst in July 2007. IGR J17098–3628 was detected for the first time by *INTEGRAL* in 2005 (while IGR J17091–3624 was not visible) and it has then remained detectable with variable flux in the soft X-ray energy range ($E < 20$ keV) up to now. On the basis of their spectral behavior, both IGR J17091–3624 and IGR J17098–3628 are classified as probable BHCs (Lutovinov & Revnivtsev 2003; Grebenev et al. 2007).

In this paper, we present the results of a long term monitoring, primarily at high energy, of these two transients, and discuss the identification in archival data of a radio counterpart for IGR J17091–3624. The date, the exposure time, the instruments used and the sources detection of the different observations are summarized in Table 1. The paper is organized as follows: in §1.1 and §1.2, we introduce the two BHCs; in §2, we describe our observations; in §3, we present the results; finally, we give a discussion and present our conclusions in §4.

1.1. IGR J17091–3624

IGR J17091–3624 was discovered by *INTEGRAL*/IBIS during a Galactic Center observation on 2003 April 14–15 (Kuulkers et al. 2003). Initially, the flux was ~ 20 mCrab in the 40–100 keV energy band exhibiting a hard spectrum, while it was not detected in the 15–40 keV band, with an upper limit of ~ 10 mCrab. During subsequent observations of the Galactic Center Deep Exposure (GCDE) on 2003 April 15–16, the source flux increased to ~ 40 mCrab in the 40–100 keV band and to 25 mCrab in the 15–40 keV (the IBIS flux statistical error is less than 10%).

Immediately after the *INTEGRAL* discovery, providing the position of IGR J17091–3624, an *RXTE* observation was performed and the source was then searched in the X-ray catalogs. IGR J17091–3624 was found in the archival data of both the TTM telescope on board the KVANT module of the *Mir* orbital station (Revnivtsev et al. 2003), and in the *BeppoSAX* WFC (in 't Zand et al. 2003). A first study of the IBIS/ISGRI spectral evolution of the source (Lutovinov & Revnivtsev 2003, 2005) showed a source hardening with a photon index changing from $\Gamma = 2.2 \pm 0.1$ to $\Gamma = 1.6 \pm 0.1$ from 2003 April to 2003 August. A

Table 1: IGR J17098-3628 and IGR J17091-3624 observations log from 2003 until September 2007

Start date	End date	Satellite	Detected sources	Exposure
2003-04-12	2003-04-21	<i>INTEGRAL</i>	IGR J17091-3624 ^a	15 ks
2003-04-20	...	<i>RXTE</i>	IGR J17091-3624 ^b	4.5 ks
2003-04-23	2003-05-09	VLA ^c	IGR J17091/IGR J17098	500 s
2003-08-10	2004-04-02	<i>INTEGRAL</i>	IGR J17091-3624 ^a	154 ks
2005-02-20	2005-05-01	<i>INTEGRAL</i>	IGR J17098-3628 ^d	300 ks
2005-03-29	...	<i>RXTE</i>	IGR J17098-3628 ^d	2.1 ks
2005-05-01	2007-02-28	<i>INTEGRAL</i>	IGR J17098-3628	450 ks
2005-05-01	2005-09-14	<i>Swift</i>	IGR J17098-3628	8.5 ks
2006-08-25	...	<i>XMM</i>	IGR J17098-3628	8 ks
2007-02-19	...	<i>XMM</i>	IGR J17098-3628	16 ks
2007-07-19	2007-08-07	<i>Swift</i>	IGR J17091/IGR J17098	11 ks
2007-08-25	2007-09-30	<i>INTEGRAL</i>	IGR J17091-3624	50 ks

^aCapitanio et al. 2006

^bLutovinov & Revnivtsev 2003

^cRupen et al. 2003

^dGrebenev et al. 2007

subsequent detailed analysis of the IBIS, JEM-X and *RXTE*/PCA data of the entire outburst duration (Capitanio et al. 2006) revealed an indication of an hysteresis like behavior and the presence of a hot disc black body emission component during the source softening.

From the investigations reported above, IGR J17091–3624 appears as a moderately bright variable transient source, with a flaring activity in 1994 October (*Mir*/KVANT/TTM), 1996 September (*BeppoSAX*/WFC), 2001 September (*BeppoSAX*/WFC, in ’t Zand et al. 2003), 2003 April (*INTEGRAL*/IBIS, Kuulkers et al. 2003), and 2007 July. In this paper, we report on the last episode of activity, as well as on limits on the quiescent state.

1.2. IGR J17098–3628

IGR J17098–3628 was detected for the first time with *INTEGRAL*/IBIS 9.4’ off IGR J17091–3624 (Grebenev et al. 2005) during deep Open Program observations of the Galactic Center region on 2005 March 24. The average fluxes were 28.2 ± 1.4 and 38.7 ± 2.8 mCrab in the 18–45 and 45–80 keV bands, respectively. Further analysis (Grebenev et al. 2005b) reported that the source was evolving in both brightness and spectral shape with an indication of softening.

From 2005 March 29 to April 4, an observational campaign was performed with *RXTE*/PCA. The spectral shape given by *INTEGRAL*/IBIS and *RXTE*/PCA varied throughout the observations and it is modelled by a soft black body emission component plus a hard tail and an absorption consistent with the Galactic one (Grebenev et al. 2007). The spectral variations suggested that this source was an X-ray nova going in outburst and probably a BHC (Grebenev et al. 2007).

The source was then observed with the *Swift* satellite (Kennea et al. 2005) with an exposure time of 2.8 ks on 2005 May 1; it was quite bright, with an estimated flux of 1.3×10^{-9} ergs s^{−1} cm^{−2} in the 0.5–10 keV energy band. The analysis of the *Swift*/XRT data for IGR J17098–3628 refined the source coordinates as follows: RA 17^h 09^m 45.9^s, Dec −36° 27′ 57″ (J2000), with an uncertainty radius of about 5″ (Kennea et al. 2005). This position is 30″ from the *INTEGRAL* position reported by Grebenev et al. (2005).

Following the soft X-ray detection of IGR J17098–3628 with *Swift*/XRT, a probable radio counterpart has also been found (Rupen et al. 2005). In particular, the first data set of four consecutive Very Large Array (VLA) radio observations, made on 2005 March 31, April 5, April 12, and May 4 at 4.86 GHz, showed only one significant radio source within the 2′ *INTEGRAL* error circle, located at RA 17^h 09^m 45.934^s $\pm$ 0.011s, Dec −36° 27′ 57.30″ $\pm$ 0.55″ (J2000) (Rupen et al. 2005).

Thanks to the radio observations, a probable optical/infrared identification has been found within the 2MASS All-Sky Catalog and the SuperCOSMOS Sky Survey (Steehgs et al. 2005b; Blustin et al. 2005; Rupen et al. 2005; Steeghs et al. 2005). On the base of the optical identification, Grebenev et al. (2007) estimated the source distance as $d = 10.5$ kpc and an upper limit on the inclination angle: $\geq 77^\circ$.

2. The multi-wavelength observation campaigns

We report here on the analyzed data collected by *INTEGRAL*, *Swift* and *XMM* from 2005 till 2007. To illustrate the need for a multi-wavelength campaign with good positional accuracy, we show in Figure 1 two *INTEGRAL* mosaic images of the region of the two BHCs. In the left panel, we show the region during the IGR J17091–3624 outburst in 2003 (IGR J17098–3628 was switched off), while in the right panel we show the same zone when only IGR J17098–3628 was visible.

The two sources are too close to each other to be resolved by X-ray and γ -ray monitors such as *RXTE*/ASM and *Swift*/BAT. Therefore, it is not possible to measure their flux separately and it is rather difficult to follow their individual activities. Observations with better positional accuracy are needed to assign each flux variation to one of the two sources with confidence. As an example, Figure 2 shows the contaminated *RXTE*/ASM IGR J17098–3628 plus IGR J17091–3624 light curve between 1.5–12 keV from 2001 until the beginning of 2008. The black and gray arrows represent the periods of *BeppoSAX*, *Swift*/XRT and *XMM* observation campaigns that verified the activity of respectively IGR J17091–3624 and IGR J17098–3628.

The X-ray Telescope (XRT) on board *Swift* satellite is an imaging instrument operating in the 0.2–10 keV energy range with a single photon point spread function of $18''$ (half-power diameter) and a spectral resolution of 140 eV at 5.9 keV. We have collected all the XRT data available from the NASA HEASARC public archive¹. The XRT data were processed using the most recently available standard *Swift* tools: XRT software version 0.11.4, FTOOLS version 6.3.1, and XSPEC version 11.3.2. The ancillary response files were generated with the *xrtpipeline* task *xrtmkarf*. The channels of each spectrum were re-binned in order to achieve a minimum of 100 counts per bin. The *Swift*/BAT transient monitor light curve was provided by the *Swift*/BAT team².

¹<http://heasarc.gsfc.nasa.gov/docs/archive.html>

²<http://swift.gsfc.nasa.gov/docs/swift/results/transients/index.html>




Fig. 1.— Left: 2003 IBIS/*INTEGRAL* 80 ks mosaic image (20-40 keV) of IGR J17091–3624 and IGR J17098–3628 zone: only IGR J17091–3624 is visible. Right: 2005 IBIS/*INTEGRAL* 120 ks mosaic image (20-40 keV) of IGR J17091–3624 and IGR J17098–3628 zone: only IGR J17098–3628 is visible




Fig. 2.— IGR J17098–3628 *RXTE*/ASM light curve contaminated by IGR J17091–3624 outbursts from 2001 till January 2008. The black triangles and gray arrows represent the periods of *BeppoSAX*, *Swift*/XRT and *XMM* observation campaigns that verified the activity of respectively IGR J17091–3624 and IGR J17098–3628. The time is expressed in Modified Julian Date (52000 (MJD)= 2001-04-01; 53500 (MJD)= 2005-05-10; 54500 (MJD)= 2008-02-04).

We also analyzed two *XMM* pointed observations performed on 2006 August 25 and 2007 February 19. *XMM* data were reprocessed using the *XMM*-Newton Standard Analysis Software (SAS) version 7.0, employing the latest available calibration files. Only single X-ray events (PATTERN=0) were taken into account for the PN; the standard selection filter FLAG=0 was applied. Exposures have been filtered for periods of high background. Since pile-up was present in both observations, source counts were extracted from annular regions of typically 90'' external radius centered on the source, and the central 7'' of the PSF have been excised; background spectra were extracted from circular regions close to the source or from source-free regions of typically 20'' of radius. The ancillary response matrices (ARFs) and the detector response matrices (RMs) were generated using the *XMM*-SAS tasks *arfgen* and *rmfgen*; spectral channels were re-binned in order to achieve a minimum of 100 counts per each bin.

The analyzed *INTEGRAL* data set consists of all Key Program and public observations from 2005 May until the end of 2007. The *INTEGRAL* data reduction of both the X-ray monitor JEM-X (Lund et al. 2003) and the γ -ray telescope IBIS (Ubertini et al. 2003) were performed using the latest release of the standard Offline Scientific Analysis (OSA) (Goldwurm et al. 2003) version 7.

We have also searched in the radio archives for data containing IGR J17091–3624 and IGR J17098–3628 observations. Very Large Array (VLA)³ observations were taken for a few epochs in 2003, following the outburst of IGR J17091–3624, and in 2005, after the IGR J17098–3628 detection. We have obtained and calibrated the data in AIPS with the standard procedures.

3. Results

3.1. Swift 2005 Observations

The 2005 *Swift* observations were performed from May 1 until September 24, after the peak of the IGR J17098–3628 outburst. A detailed analysis of this outburst peak has been reported by Grebenev et al. (2007). We report here the spectral analysis of 6 pointings (about 2 ks each) that were performed in Window Timing (WT). Figure 3 shows the image of one XRT pointing observation performed in photon counting mode in order to detect the refined position of the source; as the Figure shows, IGR J17098–3628 is quite bright while

³The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation.

IGR J17091–3624 is not visible.

During all the *Swift* observations, IGR J17098–3628 showed a slightly variable flux of about 3% with a constant spectral shape. The best fit for each individual observation is a multicolor disc black body component [*diskbb* in XSPEC Mitsuda et al. (1984)] with an internal temperature of about 0.9 keV; the equivalent hydrogen column [$N_H = (0.88 \pm 0.02) \times 10^{22} \text{ cm}^{-2}$] is consistent with the galactic one. Table 2 shows the spectral parameters of the fit for each observation.

The *INTEGRAL*/IBIS analysis of the data collected in the same period of the *Swift* observation campaign did not reveal any source emission above 15 keV up to 3σ level. The ASM monitor detected the source with an average flux of about 30 mCrab, that is a value slightly above the detection limit of JEM-X monitor (about 20–25 mCrab). Moreover, the source has been observed only in the JEM-X partially coded field of view because of dithering strategy used for *INTEGRAL* observations. For these reasons, the X-ray monitor JEM-X did not detect the source in each single science window at the 3σ level. However, adding the JEM-X science windows together yields a total exposure time of 450 ks which then provides a 36σ level detection of the source in the mosaic.

3.2. XMM 2006–2007 Observations

XMM observations of the field containing both sources were performed in 2006 August 29 and in 2007 February 19. These observations revealed that IGR J17091–3624 still did not show any detectable emission, while IGR J17098–3628 was in a relatively bright state. *INTEGRAL* observed the field containing the two sources at the same time as *XMM*, with no detections from either sources; the *INTEGRAL* upper limit flux for a single pointing (2400 s) is about 20 mCrab in the range 20–100 keV.

3.2.1. IGR J17091–3624

No detectable emission was revealed from IGR J17091–3624. Since the distance of this source is still unknown, we assumed IGR J17091–3624 as located at the Galactic Center (8 kpc). The corresponding best upper limit luminosity, derived for the quiescent state of this source and based on the *XMM* observations (exposure time 7 ks), is $L \leq 7 \times 10^{32} \text{ ergs}^{-1}$ (Watson et al. 2001).




Fig. 3.— 350 s *Swift*/XRT row image during 2005. The observation was pointed on IGR J17091–3624, which is not detected. IGR J17098–3628 is quite bright and affected by pileup.

Table 2: XRT fit parameters of the 2005 IGR J17098–3628 campaign (errors are at 90% confidence level): T_{in} = inner temperature of the accretion disk, N_{disk} = normalization constant of the *diskbb* model, $\text{Flux}_{0.5-10}$ = unabsorbed model flux between 0.5-10 keV, χ^2_{red} = reduced χ square

pointing	Esposure	T_{in}	N_{disk}	$\text{Flux}_{0.5-10}$	χ^2_{red}
Date	ks	keV	-	ergs cm ⁻² s ⁻¹	-
2005-05-01	0.4	$1.09^{0.01}_{-0.01}$	76^{4}_{-4}	2.4×10^{-9}	1.08
2005-05-10	1.0	$1.06^{0.02}_{-0.02}$	80^{6}_{-6}	1.2×10^{-9}	0.98
2005-05-21	1.0	$0.98^{0.02}_{-0.02}$	87^{8}_{-7}	1.6×10^{-9}	0.99
2005-08-13	1.5	$0.82^{0.02}_{-0.02}$	95^{11}_{-10}	9.6×10^{-10}	1.28
2005-09-05	2.1	$0.93^{0.01}_{-0.01}$	68^{5}_{-5}	1.1×10^{-9}	1.23
2005-09-14	3.2	$0.93^{0.01}_{-0.01}$	73^{4}_{-4}	1.5×10^{-9}	1.20




Fig. 4.— *XMM*/EPN image (16 ks) of the 2007-02-19 pointing observation. IGR J17091-3624 is not visible while IGR J17098-3628 is bright and affected by pile up (The dark bar in the image is due to the gap between two adjoining CCDs).

3.2.2. IGR J17098–3628

During August 2006 and February 2007, the *XMM* spectral shape of the source did not show, once more, any important variation. The only noticeable difference was the flux decrease of about 20% between the two pointings. Both spectra are characterized by an absorbed multicolor disc black body emission with an inner temperature that varied from about 1 keV in the first pointing to about 0.9 keV in the second pointing (see Table 3). The equivalent hydrogen column, $N_H = 0.8 \times 10^{22} \text{ cm}^{-2}$ is in good agreement with that measured by XRT in 2005 and with the galactic value. Table 3 shows the fit parameters of both pointings. There is no evidence in the spectral shape of a reflection component or Fe emission line according with the 2005 XRT data set analysis. Figure 5 shows the unfolded spectrum of the two *XMM*/PN pointings.

This behavior is confirmed by the *RXTE*/ASM light curve from 2005 May 1 to 2007 April 10. In this period, IGR J17091–3624 was in quiescence and the ASM light curve is only due to IGR J17098–3628 flux emission. As Figure 6 shows, the *RXTE*/ASM hardness-intensity diagram confirms that the source varied only the flux with a constant hardness ratio.

3.3. The July 2007 *Swift*/XRT monitoring: the new IGR J17091–3624 outburst

On 2007 July 9, as a consequence of an increase in the *Swift*/BAT and *RXTE*/ASM light curves, a *Swift*/XRT 2.5 ks ToO was granted. The observation was performed in window timing mode in order to avoid the pileup and revealed the presence of two sources in the XRT field of view, both of which were causing the flux increase in the *RXTE*/PCA light curve. Another 500 s pointing observation, performed in photon counting mode in order to locate the position of the two sources, revealed the presence of both IGR J17098–3628 and IGR J17091–3624, which was newly in outburst. Both sources were in relatively bright state

Table 3: PN fit parameters of the two *XMM* pointings (errors are at 90% confidence level): T_{in} = inner temperature of the accretion disk, N_{disk} = normalization constant of the *diskbb* model, $\text{Flux}_{0.5-10}$ = unabsorbed model flux between 0.5-10 keV, χ^2_{red} = reduced χ square

pointing	Exposure	T_{in}	N_{disk}	$F_{0.5-10}$	χ^2_{red}
Date	ks	keV	...	ergs $\text{cm}^{-2} \text{ s}^{-1}$	...
2006-08-25	8	$1.04^{+0.01}_{-0.01}$	47^{+3}_{-3}	1.2×10^{-9}	0.85
2007-02-19	16	$0.87^{+0.01}_{-0.01}$	65^{+2}_{-2}	8×10^{-10}	0.97




Fig. 5.— *XMM*/PN unfolded spectra of IGR J17098–3628. left panel: 2006-08-25 spectrum. Right panel: 2007-02-19 spectrum. The data are fitted with *diskbb* model. Fit parameter values are reported in Table 3




Fig. 6.— *RXTE*/ASM hardness-intensity diagram of the period from 2005 May 1 until 2007 April 10. In this period IGR J17091–3624 was in quiescence and the ASM flux emission is due only to IGR J17098–3628. The hardness ratio is defined as: $HR = (flux_B/flux_A)$ where A and B are respectively the 1.5–3 keV and 3–5 keV energy bands. The plot clearly shows that the hardness value have not changed all over the observation period.

as Figure 7 shows.

A *Swift* monitoring campaign was then requested for a total of 3 more observations, 2 ks each (one per week between 2007 July 24 and 2007 August 7) pointed on IGR J17091–3624. The key outcomes of this observation campaign for each source are summarized in the following sections.

3.3.1. IGR J17098–3628

IGR J17098–3628 has been observed during only three of the pointings associated with the observational campaign, for a total exposure of ~ 8 ks. The source flux only showed flux variations of about 30% in the range 0.5–10 keV as was inferred from the light curve shown in Figure 8. The source spectrum revealed only minor changes and the best overall fit is represented by a *diskbb* component with an inner temperature of about 0.9–1 keV.

3.3.2. IGR J17091–3624

This XRT ToO was a unique opportunity to catch the new outburst of IGR J17091–3624 from the very beginning and also provided the refined position of the source (Kennea & Capitanio 2007): RA $17^h 09^m 07.6^s$, Dec $-36^\circ 24' 24.9''$ (J2000), with an estimated uncertainty radius of $3.6''$ (90% confidence), consistent with the *INTEGRAL* position. However, this X-ray position rules out the tentative radio counterpart previously proposed for IGR J17091–3624 (Rupen et al. 2005; Pandey et al. 2006), which lies outside the XRT error circle, approximately $86''$ away. This location also rules out the optical counterpart suggested by Negueruela & Schurch (2006) which was also based on the radio counterpart. During the *Swift* observations the 0.5–10 keV flux of IGR J17091–3624 increased from 5×10^{-10} (ergs cm $^{-2}$ s $^{-1}$) to 2×10^{-9} (ergs cm $^{-2}$ s $^{-1}$). Figure 9 shows the 0.1–10 keV light curve of the 2007 XRT monitoring campaign.

In Figure 10, we show respectively the *Swift*/BAT 15–50 keV (top panel) and *RXTE*/ASM 2–10 keV (bottom panel) light curves at the time of the 2007 outburst. The luminosity peak present in both light curves is considered to be mainly due to the IGR J17091–3634 outburst, since the contamination from the IGR J17098–3628 flux should not imply such a strong variation of the flux. In fact, as discussed before, IGR J17098–3628 showed a soft spectrum and a flux variation clearly lower than the one reported by the two light curves of Figure 10.

However, the BAT and ASM light curve show a behavior typical of that expected at the beginning of a black hole binary outburst, with the hard luminosity peak preceding the




Fig. 7.— *Swift*/XRT 0.1-10 keV image of the July 2007 ToO *Swift* 500s observation. The observation has been performed in photon counting mode in order to locate the position of the sources. IGR J17098–3628 is affected by pile up.




Fig. 8.— *Swift*/XRT 0.1-10 keV light curve of IGR J17098-3628 from 2007-07-09 to 2007-07-17 (bin time: 1000 s). The gap in the light curve is due to the time interval between the second and the third XRT pointing




Fig. 9.— *Swift*/XRT (0.1-10 keV) light curve of IGR J17091-3624 from 2007-07-09 to 2007-08-07 (bin time: 1000 s).

soft one. This behavior is also confirmed by the spectral analysis: a power law component with a photon index of 1.4 ± 0.1 represents the best fit of the first two observations (9 July and 16 July 2007). For the spectra of the subsequent observations, it is necessary to add a multicolor disc black body component to the power law. Table 4 summarizes the spectral parameter values of the XRT 2007 observation campaign.

INTEGRAL detected this source only at the end of the outburst because of visibility constraints. Unfortunately the source was outside the field of view of the *INTEGRAL* X-ray monitor JEM-X, thus only the IBIS high energy data were available. Figure 11 shows the 20-100 keV IBIS image of the field of view which includes the two sources between August and September 2007 (for a total exposure time of ~ 50 ks). As Figure 11 shows IBIS detected above 20 keV only IGR J17091–3624.

The first *INTEGRAL* IGR J17091–3624 observation of the two sources field (2007-08-25) was performed 18 days after the last XRT observation (2007-08-07). However, as Figure 12 shows, XRT and IBIS spectra are in good agreement. The *INTEGRAL* observations, after the end of the XRT campaign, revealed that the spectrum of the source continued its softening with a photon index that varied from 2 to $\simeq 3$ at the end of September 2007.

3.4. Radio Counterparts

A plot of the NVSS data in the region of IGR J17091–3624 and IGR J17098–3628 is shown in Figure 13 together with the position and associated uncertainties of the *INTEGRAL* and *Swift*/XRT measurements (marked by vertical and diagonal crosses, respectively) and the tentative radio counterparts from VLA literature data (Rupen et al. 2003, 2005). In the region of IGR J17098–3628, there is clearly an excellent agreement between the radio and X-ray positions, which gives a high confidence identification. The NVSS contours themselves show some emission in this region, although it is quite extended and only significant at the 2.5σ level. In the region of IGR 17091–3624, the refined *Swift*/XRT position is clearly not consistent with the proposed radio counterpart. No signal is detected in the NVSS image in that region.

We have therefore searched the VLA archive for higher resolution images of these regions, primarily to find a radio counterpart to IGR J17091–3624 but also in order to study the time and the spectral evolution of the sources. We have found archival data from 2003 in the region of IGR J17091–3624 and from 2005 in the region of IGR J17098–3628. The data from 2003 were taken in the compact D configuration on Apr 23, Apr 26, May 06, and May 09, with exposures of a few minutes. We show an 8.4 GHz image of the IGR J17091–3624 region in




Fig. 10.— Top panel: IGR J17091–3624 *Swift*/BAT (15–50 keV) light curve. Bottom panel: IGR J17091–3624 *RXTE*/ASM (2–10 keV). Both curves are contaminated by the IGR J17098–3628 flux emission. The time is expressed in Modified Julian Date (54000 (MJD)= 2006-09-22; 54200 (MJD)= 2007-04-10; 54400 (MJD)= 2007-10-27)

Table 4: The fit parameters of the observed spectra of IGR J17091–3624 during 2007 *Swift*/XRT monitor campaign (errors are at 90% confidence level): T_{in} = disk inner temperature, N_{disk} = *diskbb* model normalization constant, Γ = power law photon index, N_{pow} = power law model normalization constant, $Flux_{0.5-10}$ = unabsorbed model flux between 0.5-10 keV, χ^2_{red} = reduced χ square.

pointing	Exposure	T_{in}	N_{disk}	Γ	N_{pow}	$Flux_{0.5-10}$	χ^2_{red}
Date	ks	keV	...	...	pho keV ⁻¹ cm ⁻² s ⁻¹	ergs cm ⁻² s ⁻¹	...
2007-07-09	2.4	...	...	$1.4^{0.1}_{-0.1}$	$0.05^{0.01}_{-0.01}$	4.7×10^{-10}	0.96
2007-07-16	3.4	...	...	$1.3^{0.1}_{-0.1}$	$0.028^{0.003}_{-0.003}$	3.0×10^{-10}	1.10
2007-07-24	2.6	$0.86^{0.03}_{-0.03}$	69^{13}_{-13}	$1.6^{0.1}_{-0.1}$	$0.12^{0.02}_{-0.02}$	1.8×10^{-9}	1.20
2007-08-07	1.9	$0.75^{0.05}_{-0.07}$	57^{30}_{-23}	$2.0^{0.1}_{-0.1}$	$0.23^{0.04}_{-0.03}$	2.1×10^{-9}	1.08




Fig. 11.— IGR J17091–3624 *INTEGRAL*/IBIS 20-100 keV image between August and September 2007 (total exposure time of ~ 50 ks).

Figure 14, superimposed to the *Swift*/XRT position. A radio source is clearly revealed in a position consistent with the high energy error circle. This component is present in the three epochs at 8.4 GHz, and shows an increase in total flux. Data at 5 GHz are also available at three epochs; the source is marginally detected in the first one (3σ , 5 GHz data only), and clearly revealed in the next two. For the epochs where simultaneous data are available, a comparison to the 8.4 GHz flux reveals an inverted spectrum. Table 5 reports on the measured flux densities; the position of the radio source is RA $17^h 09^m 07.5^s \pm 0.5^s$, Dec $-36^\circ 24' 24'' \pm 3''$.

4. Discussion and Conclusions

The huge amount of data collected and the broad energy coverage allowed us to perform a detailed spectral analysis of these two sources, following their evolution during more than two years and fixing upper limits when one of the sources was not visible. In the following a detailed summary of the results for both sources can be found.

4.1. IGR J17098–3628

Our analysis, spanning from May 2005 to September 2007, proves that the source spectrum shows a soft black body component with an internal temperature of about ~ 1 keV and an internal radius comparable to the last stable orbit of the accretion disc. The luminosity due to soft component (assuming $d = 10.5$ kpc, Grebenev et al. 2007) varied during the two years of our observational campaign from $\sim 2 \times 10^{37}$ to $\sim 5 \times 10^{36}$ ergs s $^{-1}$. The flux variation range and the spectral parameters are comparable with the ones reported by

Table 5: Flux density of the radio counterpart of IGR J17091–3624. The observation of 2003 Apr 23 was at 5 GHz only and is a 3σ detection; 2003 May 09 is only at 8.4 GHz. Spectral index is defined according to $S(\nu) \propto \nu^{-\alpha}$.

Date	$S_{5\text{ GHz}}$	$S_{8.4\text{ GHz}}$	Spectral index
-	(mJy)	(mJy)	-
2003 Apr 23	0.32	...	...
2003 Apr 26	0.48	0.56	−0.3
2003 May 06	0.42	0.67	−0.8
2003 May 09	...	0.73	...

Grebenev et al. (2007) during the first phases of the outburst on April 2005. In our data set each flux variation seems not to be correlated with any relevant spectral features, as confirmed by the lack of evident variation of the *RXTE*/ASM hardness-intensity diagram (Figure 6). However respect to the previous April 2005 observations, we did not detect any power law emission at higher *INTEGRAL*/IBIS energy band. This indicates that the spectral shape was evolved from the first phase of the outburst being dominated only by a thermal disc component coming from the accretion disc around the compact object.

We can conclude that this source seems to have spent 2.5 years in the high soft state with a disc black body component substantially identical to the one previously observed at the beginning of the outburst (Grebenev et al. 2007). Curiously the only difference is the lack of any high energy emission. In fact the hard component fell below the detection limit of IBIS after about three months from the beginning of the outburst. Hence the geometry and the temperature of the accretion disc have not shown any significant variation up to now, on the other end the power law emission quenched probably because of the electron temperature of the corona fallen below the disc seed photons temperature making the inverse Compton scattering processes inefficient.

4.2. IGR J17091–3624

Previous studies have demonstrated that the X-ray luminosity of the BHCs in quiescence is lower than that of neutron star X-ray binaries and falls below $\sim 10^{32}$ – 10^{33} ergs s $^{-1}$ (Campana et al. 2001). In the quiescent state, neutron star X-ray binaries are normally detected and hence the *XMM* upper limit of IGR J17091–3624 is a good indication that the source is a BHC.

The *Swift* ToO caught the source at the beginning of its outburst when it was still in a low/hard state. The best fit of the first observation (2007-07-09/16) is a power law spectrum with $\Gamma=1.4\pm 0.1$. The source spectrum substantially softened and the black body component became dominant with an increasing temperature and a steeper power law. *INTEGRAL* continued to monitor the source with IBIS after the end of the XRT observational campaign, up to the end of September 2007. These observations confirm the softening of the source. A power law component without an energy cutoff provided the best fit of the IBIS spectra between 20 and 80 keV, probably due to jets or corona reprocessing of disc seed photons. The power law photon index varied during two month period from about 1.4 (beginning of July) to about 3 (end of September). Unfortunately the relatively low source flux at high energies (7×10^{-10} ergs cm $^{-2}$ s $^{-1}$ between 20-100 keV) and the variation of the spectral shape did not provide sufficient statistics to extend the spectra up to 80 keV and to verify

the presence or not of an high energy cutoff in the spectrum.

Thanks to the XRT refined position, we found the radio counterpart of the source. As soon as 9 days after the detection by IBIS in 2003, a radio source at the sub-mJy level was detected at 5 GHz. The source has been detected also at 8.4 GHz, and it showed an increase in flux over the subsequent two weeks. The spectrum is inverted, characteristic of self-absorbed synchrotron radiation from a compact jet. This behavior is typical of BHC in low hard states. Although it is difficult to guess the spectral shape at higher frequencies, it is reasonable to estimate that the total radiative luminosity of the compact jet is of the order of 10^{31} erg sec $^{-1}$, assuming the distance to the Galactic Center. No information is available after the 2007 outburst, and it will be clearly valuable to obtain new radio observations after future episodes of activity.

Chaty et al. 2008, on the basis of ESO NTT observations, report on two possible infrared counterparts of IGR J17091–3624 consistent with the *Swift*/XRT error box. Our refined position, based on the radio observations, could support the identification of the real infrared counterpart of this source.

Little is known about nature, duration and recurrence of outbursts in transient X-ray binaries. It is probable that the outbursts are due to randomly acting factors such as the mass transfer variations or truncation of the inner disc radius. However for IGR J17091–3624 five outbursts are known from 1994 to 2007. Thus, it appears that this source goes in outburst every three or five years. Generally the flux varies from 5 mCrab to about 20 mCrab in the range 2-10 keV over a period of few months. The 2003 outburst was the first detected at higher energy range (20-150 keV) for one year period. Comparing the two hard states of respectively the 2003 and 2007 outbursts, the last one seems to be the hardest. In fact it was possible to determine the high energy power law cutoff of the 2003 hard state ($\sim$ 49 keV, Capitanio et al. 2006) while in the 2007 hard state the cutoff is not detectable up to 80 keV. Concerning the soft state the 2003 outburst has an higher temperature black body emission (Capitanio et al. 2006) that does not appear to be present in the 2007 outburst. However the XRT monitoring campaign observed the source only at the beginning of its transition to the soft state and unfortunately in the subsequent *INTEGRAL* monitoring the source was outside the JEM-X field of view and so it was not possible to follow the entire evolution of the black body temperature.

This work has been supported by the Italian Space Agency through grants I/008/07/0 and I/088/06/0. We acknowledge the use of public data from the *Swift* data archive and all the *Swift* team for its support and Memmo Federici for supervising the *INTEGRAL* data analysis. Particular thanks are due to Erick Kuulkeers for making his private data available

for analysis.

REFERENCES

- Bird, A. J., Malizia, A., Bazzano, A. et al., 2007, ApJS, 170, 175,
- Blustin, A. J., Kong, A. K. H., Wu, K. et al., Atel 479, 2005
- Campana, S., A. N., Parmar and Stella, L. A&A, 372, 241, 2001
- Capitanio, F., Bazzano, A., Ubertini, P. et al. ApJ, 643, 376, 2006
- Chaty, S., Rahoui, F., Foellmi, F. et al. A&A in press 2008 (astro-ph: 0802.1774)
- Goldwrum, A., Daid, P., Foschini, L. et al. A&A, 411, 1223, 2003
- Grebnev, S., A., Molkov, S., V., Revnivitsev, M., G. & Sunyaev, 2007, ESA SP, 622, 373
(astro-ph: 0709.2313)
- Grebenev, S. A., Molkov, S. V., Sunyaev, R. A. 2005, Atel 444
- Grebenev, S. A., Molkov, S. V., Revnivitsev, M. G. & Sunyaev, R. A., 2005 Atel 447
- Kennea, J. A. & Capitanio, F. 2007, Atel 1140
- Kennea, J. A., Burrows, D. N., Nousek, J. A. et al., 2005, Atel 476.
- Kuulkers, E., Lutovinov, A., Parmar, a., Capitanio, F. et al., 2003 Atel 149
- Kong, A. K. H. 2005, Atel 477
- Lund, N., Budz-Joergensen, C., Westgaard, N. J., 2003 A&A, 411, L231
- Lutovinov, A. & Revnivitsev, M. 2003, Astr. Lett., 29, 719
- Lutovinov, A., Revnivitsev, M., Molkov, S. and Sunyaev, R. 2005, A.&A., 430, 997
- Mitsuda, K., Inoue, H., Koyama et al., 1984, PASJ, 36, 741
- Mowlavi, N., Kuulkers, E., Rodriguez et al., 2005, Atel 453
- Negueruela, I. & Schurch, M. P. E., 2006, A&A, 461, 631
- in 't Zand, J. J. M., Heise, J., Lowes, P. & Ubertini, P. 2003, ATel, 160

- Pandey, M., Manchanda, R. K., Rao, A. P., Durouchoux, P. & Ishwara-Chandra 2006
- Revnivtsev, M., Gilfanov, M., Churazov, E., & Sunyaev R. 2003, ATel, 150
- Rupen, M. P., Mioduszewski, A. J., & Dhawan, V. 2003, ATel, 152
- Rupen, M. P., Mioduszewski, A. J., Dhawan, V., 2005, Atel 490
- Steeghs, D., Torres, M. A. P., Jonker, P. G. et al., 2005, Atel 494
- Steeghs, D., Torres, M. A. P., Jonker et al., 2005, Atel 478
- Ubertini, P., Lebrun, F., Di Cocco, G. et al., A&A, 411, L131, 2003
- Watson, M. G., Augures, J.-L., Ballet, J. et al., A&A 364, L51, 2001




Fig. 12.— IGR J17091–3624 2007 July outburst: XRT-IBIS combined spectrum: the XRT observation (2 ks) had been performed 18 days before that of IBIS (14 ks). The best fit model is represented by a multicolor disc black body (red dashed line) plus a power law component (green dashed line). The total model is represented in the figure by the solid green line. The spectral parameters are: $T_{in} = 0.75 \pm 0.1 \text{ keV}$, $N_{disk} = 57 \pm 25$, $\Gamma = 2 \pm 0.1$. The flux between 0.5–100 keV is : $9 \times 10^{-9} \text{ ergs cm}^{-2} \text{ s}^{-1}$




Fig. 13.— NVSS image of the region of IGR J17091–3624 and IGR 17098–3628, contours traced at $(-1, 1, 2, 4, \dots) \times 1 \text{ mJy beam}^{-1}$ (thin grey lines). Darker bold symbols represent IBIS positions (big plus signs), *Swift*/XRT positions (circles, 90% confidence), and previously proposed radio counterparts (crosses). The key in the bottom right corner shows the radio beam size ($45'' \times 45''$ HPBM).




Fig. 14.— Very Large Array 8.4 GHz image of the region of IGR 17091–3624, taken on 2003 May 09. Contours are traced at $(-0.3, 0.3, 0.6, 1.2)$ mJy beam⁻¹; the gray scale range is 0.2 to 1.1 mJy beam⁻¹. The circle shows the refined *Swift* position (90% confidence), and the cross shows the previously proposed radio counterpart. The key in the bottom left corner shows the radio beam size ($40.4'' \times 7.3''$ HPBM in PA 30° , measured north to east).